

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/radcr

Case Report

The subtle scaphoid fracture in childhood: A case report with literature insights ☆,☆☆

Amr Khaled, MD^a, Fathi Milhem, MD^a, Ameer awashra, MD^a, Mazen Abdalla, MD^{a,b,*},
 Mohammad M. Jaber, MD^{a,b}, Ayman alKhawaja, MD^{a,b}

^aDepartment of Medicine, An-Najah National University, Nablus, Palestine

^bDepartment of Orthopedic, An-Najah National University Hospital, Nablus, Palestine

ARTICLE INFO

Article history:

Received 25 September 2024

Revised 6 October 2024

Accepted 14 October 2024

Keywords:

Pediatric scaphoid fracture

Conservative management

ABSTRACT

We describe a 9-year-old male who suffered a linear, non-displaced scaphoid fracture following a pedestrian motor vehicle accident. Initially, X-ray imaging failed to detect the fracture, but MRI confirmed the diagnosis, highlighting the challenges in identifying scaphoid fractures in pediatric patients due to the bone's cartilaginous nature in this age group. The patient was treated with immobilization using a Plaster of Paris cast, consistent with standard management for non-displaced fractures in children. He experienced pain improvement and achieved bone union after eight weeks, returning to normal activities without complications. The discussion emphasizes the rarity of scaphoid fractures in children under 10 and underscores the increasing incidence of such injuries in pediatric populations due to sports participation. A comparative analysis of similar cases in children of different ages is included, demonstrating variations in fracture types, treatment approaches, and outcomes, providing valuable insights into the management of pediatric scaphoid fractures.

© 2024 The Authors. Published by Elsevier Inc. on behalf of University of Washington.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Introduction

Carpal fractures are more common in adults, with scaphoid fractures being the most well-known. However, due to the early participation of children in high-contact sports, the incidence of scaphoid waist fractures is rising among this younger population [1,2].

Scaphoid fractures are very rare fractures in childhood, it makes about 3% of fractures of hand and wrist, and they are

more common in male because male engage in high-impact activities and delayed growth and ossification process exposes scaphoid to injury [3,4].

Scaphoid waist fractures are most frequently observed in skeletally immature children aged 12 to 15, as this period coincides with the ossification of the scaphoid bone [5]. The unique anatomy and vascular supply of the scaphoid make these fractures particularly challenging to diagnose, even for experts. However, in children under 10 years old, these fractures are rare because the scaphoid bone is primar-

☆ Competing Interests: The authors state that they have no conflict of interest to be mentioned.

☆☆ Acknowledgment: We express our deep gratitude to the surgical ward staff for their invaluable support in completing this report.

* Corresponding author.

E-mail address: mazen.Abdalla@najah.edu (M. Abdalla).

<https://doi.org/10.1016/j.radcr.2024.10.087>

1930-0433/© 2024 The Authors. Published by Elsevier Inc. on behalf of University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

ily cartilaginous at this stage and has not yet fully ossified [6].

In this case, a 9-year-old patient was suspected of having a scaphoid fracture based on the clinical presentation, leading to difficulties and delays in diagnosis.

Case presentation

A 9-year-old male was admitted after being struck by a car in a pedestrian motor vehicle accident (MVC). He was hit from the side, causing him to fall and slide on the right side of his body.

Upon arrival at the emergency department, the patient was conscious, alert, and oriented with a Glasgow Coma Scale (GCS) score of 15/15. And the patient was stable after primary and secondary survey. The patient was complaining of right wrist pain. Examination of the right wrist showed mild swelling over the lateral aspect of the wrist and tenderness over the anatomical snuffbox, though no deformity was noted. A right wrist X-ray did not reveal any fractures (Figs. 1 and 2).

Given the mechanism of injury and the patient's localized wrist pain, further evaluation was warranted. An orthopedic consultation was obtained, and an MRI of the right wrist showed findings consistent with a linear non displaced scaphoid fracture (Figs. 3 and 4). A Plaster of Paris cast was applied for immobilization.

During the hospital stay, the patient's vital signs remained stable, and his pain improved following analgesia and application of the cast. He was re-evaluated and was in good condition and discharged home with the cast in place. Follow-up was arranged to monitor the scaphoid fracture.



Fig. 1 – An anterior view of right wrist x-ray shows no fracture.



Fig. 2 – A lateral view of right wrist x-ray shows no fracture.



Fig. 3 – A Saggital view of the right wrist MRI, T1-weighted imaging (T1W1), reveals: A hypointense line suggestive of a scaphoid fracture (red arrows), along with edema localized to the distal scaphoid region (black arrow).



Fig. 4 – A coronal view of the right wrist MRI, using Short Tau Inversion Recovery (STIR), demonstrates: A hyperintense signal within the distal portion of the scaphoid bone, indicative of bone marrow edema (red arrow). Additionally, mild joint effusion is observed (black arrow).

Discussion

A scaphoid fracture is one of the most common wrist fractures in the pediatric population [7]. According to a retrospective analysis by Ahmed I et al., the annual incidence of scaphoid fractures is estimated to be 11 per 100,000. The most affected

age group is between 12 and 15 years, while boys under 10 years old are rarely affected [5]. However, the incidence of scaphoid fractures in pediatric patients is increasing due to greater participation in sports, demographic shifts, and advancements in diagnostic and treatment methods [8].

The ossification of the first metacarpal bone occurs simultaneously with that of the scaphoid, making it a potential marker for scaphoid maturity. The ossification of the scaphoid progresses in a distal-to-proximal direction, beginning around 5 to 6 years of age and completing by approximately 15 years and 3 months in boys, and 13 years and 4 months in girls [9]. During this period, the scaphoid remains largely cartilaginous, which not only reduces the likelihood of fractures but also complicates the diagnostic process in children [6].

This makes diagnosis with X-ray very difficult in this pediatric group, explaining the high incidence of missed scaphoid fractures, especially when associated with concurrent wrist or elbow injuries [6]. Hence, MRI is the preferred diagnostic modality in this age group, with a negative predictive value reaching 100% [10]. Although this primarily applies to the 12–15 year age group, as they are the most affected, literature regarding children under 10 years of age remains scarce [6]. This aligns with findings in the current case, where the initial wrist X-ray failed to reveal a fracture, a common occurrence. MRI, being the most sensitive tool for detecting scaphoid fractures, becomes crucial when clinical suspicion persists despite normal X-ray results.

The treatment of pediatric scaphoid fractures typically involves immobilization, with surgery being rare and generally reserved for cases of nonunion or displaced fractures [11]. In this case, the patient was treated conservatively with immobilization using a Plaster of Paris cast, which is consistent with standard treatment protocols for nondisplaced scaphoid fractures in children. Studies show that pediatric patients generally have a good prognosis with conservative management due to their high healing potential, and complications such as nonunion or avascular necrosis are uncommon [7,11].

Table 1 – Comprehensive review of multiple cases comparable to ours.

Case description	Fracture type	Treatment	Outcome	Complications
9-year-old male with scaphoid fracture after a motor vehicle accident (Our Case)	Linear scaphoid fracture	Plaster of Paris cast	Bone union after 8 weeks, returned to usual activities.	None reported
8-year-old male with scaphoid fracture after a fall [12]	Scaphoid fracture (initially undiagnosed)	Cast, delayed surgery with bone graft	Surgical union at age 16 after 7 years of follow-up	Avascular necrosis, nonunion
11-year-old male gymnast with stress fracture of the ossification center of the scaphoid [13]	Stress fracture of the ossification center	Short arm cast for 4 weeks	Bone union after 8 weeks, returned to sports after 4 months	None reported
11-year-old male with scaphoid fracture after sports injury [14]	Scaphoid fracture	Surgery with K-wire fixation and bone graft	Complete union after 10 months, full range of motion restored	Nonunion
14-year-old male with scaphoid fracture from fall during sports [2]	Proximal pole scaphoid fracture (nonunion)	Surgery with vascularized bone graft and internal fixation	Complete union after 6 months, returned to sports without restrictions	None reported

We conducted a comprehensive review of multiple pediatric scaphoid fracture cases, presented in a comparative table that highlights variations in fracture types, treatment approaches, and clinical outcomes. This detailed comparison provides a deeper understanding of the effectiveness of different treatment modalities and the potential complications associated with scaphoid fractures in children. The table serves as a valuable reference, offering insights into the diverse presentations and management strategies for this injury in the pediatric population (see Table 1).

Patient consent

- (1) Ethics approval Our institution does not require ethical approval for reporting individual cases or case series.
- (2) Written informed consent was obtained from the patient Parents for their anonymized information to be published in this article

REFERENCES

- [1] Christodoulou AG, Colton CL. Scaphoid fractures in children. *J Pediatr Orthop* 1986;6(1):37–9.
- [2] Bhatti AN, Griffin SJ, Wenham SJ. Deceptive appearance of a normal variant of scaphoid bone in a teenage patient: a diagnostic challenge. *Orthop Rev (Pavia)* 2012;4(1):e6.
- [3] Nguyen JC, Nguyen MK, Arkader A, Guariento A, Sze A, Moore ZR, et al. Age-dependent changes in pediatric scaphoid fracture pattern on radiographs. *Skeletal Radiol* 2020;49(12):2011–18.
- [4] Gause TM, Moran TE, Carr JB, Deal DN. Scaphoid waist nonunion in an 8-year-old: a rare occurrence. *Case Rep Orthop* 2019;2019:4701585.
- [5] Ahmed I, Ashton F, Tay WK, Porter D. The pediatric fracture of the scaphoid in patients aged 13 years and under: an epidemiological study. *Journal of Pediatric Orthopaedics* 2014;34(2):150.
- [6] Tataria R, Ting J, Jester A, Lindau TR, Oestreich K. Scaphoid fractures below the age of 10: case series and review of the literature. *J Wrist Surg* 2020;9(6):458–64.
- [7] Ting B, Sesko Bauer A, Abzug JM, Cornwall R, Wyrick TO, Bae DS. Pediatric scaphoid fractures. *Instr Course Lect* 2017;66:429–36.
- [8] Changing patterns of scaphoid fractures in adolescents - PubMed. [accessed 09.13.24]. <https://pubmed.ncbi.nlm.nih.gov/8199938/>.
- [9] Huckstadt T, Klitscher D, Weltzien A, Müller LP, Rommens PM, Schier F. Pediatric fractures of the carpal scaphoid: a retrospective clinical and radiological study. *J Pediatr Orthop* 2007;27(4):447–50.
- [10] Cook PA, Yu JS, Wiand W, Cook AJ, Coleman CR, Cook AJ. Suspected scaphoid fractures in skeletally immature patients: application of MRI. *J Comput Assist Tomogr* 1997;21(4):511–15.
- [11] McVeigh KH, Berger TG, Cudahy R, Dekker TM, Brigham TJ, Braxton JC. An evidence-based approach to casting and orthosis management of the pediatric, adolescent, and young adult population for injuries of the upper extremity: a review article. *Clin J Sport Med* 2021;31(2):151–62.
- [12] Davies PSE, Cox QGN. Avascular necrosis and fracture of the scaphoid in a child followed by non-union. *J Hand Surg Eur Vol.* 2019;44(1):109–10.
- [13] Fujioka H, Nishikawa T, Takagi Y, Oi T, Yoshiya S. Stress fracture of the ossification center of the scaphoid in a skeletally immature gymnast. *J Hand Surg Asian Pac Vol.* 2019;24(3):386–8.
- [14] Hamdi MF, Khelifi A. Operative management of nonunion scaphoid fracture in children: a case report and literature review. *Musculoskelet Surg* 2011;95(1):49–52.