



Identification of Pharmacophore Site of Armed Quinoxaline, Benzothiazole and Benzothiazine: POM Theoretical Screening and Experimental Evaluation of Antifungal and Antibacterial Activity

TAIBI BEN HADDA^{1*}, FATIM-ZAHRA. KHARDLI¹, MOSTAFA MIMOUNI¹, MARIA DAOUDI², ABDELALI KERBAL², BRAHIM BENNANI², ISMAIL WARAD³, HÉCTOR SALGADO-ZAMORA⁴, TEFFAHA FERGOUG⁵

¹Laboratoire de Chimie des Matériaux, Faculté des Sciences, 60000 Oujda, Morocco.

²Laboratoire Chimie Organique, Faculté des Sciences Dhar Mehraz, 30000 Fès, Morocco.

³ Department of Chemistry, Science College. King Saud University, P. O. Box 2455, Riyadh 11451, Saudi Arabia

⁴Departamento de Química Orgánica, Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional, Col. Santo Tomás 11340, México, D. F.

⁵Laboratoire de Chimie Physique des Biomolécules et Interfaces biologiques, Faculté des Sciences de la Nature et de la Vie, Université de Mascara, Mascara 29000, Algérie.

Received 7 March 2013; revised 17 April 2013; accepted 27 April 2013

Available online 7 May 2013

Abstract- A series of four different armed heterocyclic candidates; the 1-(2-methyl-2,3-dihydro-1,3-benzothiazol-2-yl)acetone (**2**), 1-(3-methyl-4*H*-1,4-benzothiazin-2-yl)ethanone (**3**), 2-[(2-aminophenyl)dithio]aniline (**4**) and 3-hydroxy-3-methyl-4-(3-methyl-2-quinoxaliny)-2-butanone (**5**) have been prepared and their antibacterial and antifungal activity were evaluated.
